

Quark-hadron duality and production of charmonia and doubly charmed baryons in e^+e^- annihilations

A. V. Berezhnoy^{1,*} and A. K. Likhoded^{2,†}

¹*SINP MSU, Moscow, Russia*

²*IHEP, Protvino, Russia*

In the framework of quark-hadron duality, the cross section values of charmonium and doubly charmed baryon production in e^+e^- -annihilations have been estimated at interaction energy $\sqrt{s} = 10.6$ GeV. It has been shown that the approach under discussion allows to resolve some contradictions between the experimental data and pQCD predictions.

I. INTRODUCTION

Until recent times, the calculation of J/ψ production cross section was done within the so-called δ -approximation, which is based on the factorization theorem of QCD. In the approach under discussion, relative momenta of valence quarks of J/ψ -meson are not taken into account in the hard part of amplitude. One supposes that both valence quarks are on-shell and each of them carries a half of the total J/ψ -meson momentum. Nevertheless, the recent experimental results of the BELLE and BABAR Collaborations on J/ψ -production in e^+e^- annihilations demand an essential revision of the calculation techniques based on QCD factorization theorem. Indeed, the cross section value of charmonium pair production in e^+e^- annihilations estimated in the framework of the δ -approximation [1] underestimates the experimental data by an order of magnitude [2, 3]. The detailed analysis has shown that such underestimation is caused by large fixed virtualities of the intermediate quark and gluon $q^2 \sim \frac{Q^2}{4}$ (Q^2 is a virtuality of the initial photon, see Fig. 1), which occur within the δ -approximation technique.

Taking into account the relative motion of valence quarks allows to remove contradictions

*Electronic address: aber@ttk.ru

†Electronic address: likhoded@mx.ihep.su

between theory and experiment, at least in principle. Calculations of the pair quarkonium production in e^+e^- annihilations, obtained within a light cone formalism [4] are in a qualitative agreement with the experimental data of the BELLE and BABAR Collaborations. In these works, the δ -shaped wave function of quarkonium was replaced by a function $f(x)$, which is "spread" on x , where x is a momentum fraction of the quarkonium carried by a valence quark in the infinite momentum frame. Thus, the longitudinal "internal motion" of the valence quarks is taken into account ¹. This spreading leads to a decrease of effective virtuality of the intermediate quark and gluon, and, therefore, to an increase of the predicted cross section value.

In spite of certain success in the description of charmonium pair production, uncertainties of the concrete model realization remain. These are uncertainties of wave function value, final c -quark virtualities, and standart uncertainties of running constant value α_s , c -quark mass value m_c , etc.

The cross section of inclusive charmonium production in e^+e^- annihilations ($e^+e^- \rightarrow \text{charmonium} + c + \bar{c}$) depends on model uncertainties to a lesser extend [6, 7], because the largest cross section fraction for such process is determined by the smallest virtualities of the intermediate quark and gluon (Fig. 1). Taking into account the relative motion of quarks changes the the result less than 10% [8], and a general dependence on choice of m_c and α_s values remains.

In this work we perform a new analysis of four charmed quark production in e^+e^- annihilations within the quark-hadron duality.

II. EXCLUSIVE AND INCLUSIVE CHARMONIUM PRODUCTION

It is reasonable to describe the cross section of four c -quark production in the process

$$e^+e^- \rightarrow c\bar{c}c\bar{c}, \quad (1)$$

in the framework of perturbative QCD. A value of $m_c = 1.25$ GeV can be chosen for the c -quark mass. This value is the same as that adopted for QCD sum rules. The strong

¹ It is worth to note that the problem of internal motion of valence quarks inside J/ψ -meson was discussed a rather long time ago. See, for example, works [5], where it was suggested to use the structure function of J/ψ in form $F_c = x^{2.2}(1-x)^{2.45}$.

coupling constant at the scale of m_c is put to $\alpha_s = 0.24$. It is worth to mention that the choice of relatively small value of c -quark mass makes a use of quark-hadron duality more convincing, because the masses of observed doubly heavy hadrons lie approximately in the center of duality interval at this m_c value.

The cross section value of process (1) can be considered as a sum of cross section values of all process with four c -quarks. Leading order calculations with the chosen values of α_s and m_c at $\sqrt{s} = 10.6$ GeV give the following cross section value of the process

$$\sigma(e^+e^- \rightarrow c\bar{c}c\bar{c}) = 372 \text{ fb.} \quad (2)$$

The cross section of pure electro-dynamical production of four c -quarks is only $1 \div 2\%$ of the above value.

There are uncertainties in prediction for the process (1) due to the choice of m_c and α_s values. The m_c and α_s used here lead to the most optimistic prediction for the cross section value of four c -quark production. This value can be considered as an upper limit of four c -quark yield. The performed estimation (2) is in accordance with the cross section value at Z^0 -boson peak [9].

As it was noted earlier [10], for the processes

$$e^+e^- \rightarrow J/\psi + c + \bar{c}, \quad (3)$$

$$e^+e^- \rightarrow \eta_c + c + \bar{c}, \quad (4)$$

the total cross section values estimated within the δ -approximation [6, 7] agree with the cross section value of production of color singlet $c\bar{c}$ -pair calculated for a some duality interval.

Using values of wave function at origin $|R_s(0)|^2 = 0.67 \text{ GeV}^3$ and the charmonium mass value 3 GeV, the following results have been obtained for J/ψ and η_c production at $\sqrt{s} = 10.6$ GeV:

$$\sigma(e^+e^- \rightarrow J/\psi + c + \bar{c}) = 104 \text{ fb}, \quad (5)$$

$$\sigma(e^+e^- \rightarrow \eta_c + c + \bar{c}) = 40 \text{ fb.} \quad (6)$$

Therefore,

$$\sigma(J/\psi)/\sigma(\eta_c) = 2.6. \quad (7)$$

A contribution of higher S - and P -wave charmonium excitations into the inclusive production cross section is about 50% [7], consequently the total cross section value can be

estimated² as 216 fb. This value can be compared to the cross section value obtained in the appropriate duality interval of mass spectrum of color singlet $c\bar{c}$ -pairs produced in the process (1):

$$2m_c < m_{c\bar{c}} < 2m_D + \Delta, \quad (8)$$

where $m_c = 1.25$ GeV, $\Delta = 0.5$ GeV, and m_D is the experimental D -meson mass. Thus, one can obtain:

$$\int_{2m_c}^{2m_D+\Delta} \frac{d\sigma(e^+e^- \rightarrow (c\bar{c})_{\text{singlet}} + c + \bar{c})}{dm_{c\bar{c}}} dm_{c\bar{c}} \Big|_{\Delta=0.5 \text{ GeV}} = 280 \text{ fb.} \quad (9)$$

This value should be compared with 216 fb. Therefore, for the case of inclusive charmonium production (3) and (4) for some Δ a fair agreement between the results of δ -approximation and the quark-hadron duality hypothesis can be achieved. For the color singlet pairs with a definite spin value ($S = 0$ or $S = 1$) one can obtain:

$$\int_{2m_c}^{2m_D+\Delta} \frac{d\sigma(e^+e^- \rightarrow (c\bar{c})_{\text{singlet}}^S + c + \bar{c})}{dm_{c\bar{c}}} dm_{c\bar{c}} \Big|_{\Delta=0.5 \text{ GeV}}^{S=1} = 204 \text{ fb,} \quad (10)$$

$$\int_{2m_c}^{2m_D+\Delta} \frac{d\sigma(e^+e^- \rightarrow (c\bar{c})_{\text{singlet}}^S + c + \bar{c})}{dm_{c\bar{c}}} dm_{c\bar{c}} \Big|_{\Delta=0.5 \text{ GeV}}^{S=0} = 76 \text{ fb.} \quad (11)$$

If one supposes that the spin-triplet $c\bar{c}$ -pair ($S = 1$) most probably transforms into a J/ψ meson and the spin singlet $c\bar{c}$ -pair ($S = 0$) most probably transforms into η_c , then

$$\sigma(J/\psi)/\sigma(\eta_c) = 2.7. \quad (12)$$

This ratio is in a good agreement with one calculated in the frame work of the δ -approximation.

Thus for the case of inclusive charmonium production, the both approaches under discussion agree. Contrary, for a case of exclusive charmonium pair production the cross section value calculated within the quark-hadron duality hypothesis is larger than the δ -approximation prediction by one order of magnitude. Such a difference occurs because the quark-hadron duality allows to take into account the internal motion of quarks inside charmonium. Indeed, a quark and an antiquark produced in duality interval (8) and fused into charmonium according to the quark-hadron duality hypothesis can have different velocities.

² Here and below the process of one-photon annihilation is discussed. Two-photon annihilation $e^+e^- \rightarrow \gamma^*\gamma^* \rightarrow J/\psi + c + \bar{c}$ contributes less than 10% into the total cross section value [11].

This decreases the effective virtuality of the intermediate gluon and quark, and increases consequently the predicted cross section value.

The exclusive charmonium pair production cross section can be estimated by picking out the both $c\bar{c}$ -pairs produced in the process (1) in appropriate duality intervals and in color-singlet state. In our calculations, we will also distinguish spin states of the pairs (S_1 and S_2 , correspondingly). The calculation results are as follows:

$$\iint_{2m_c}^{2m_D+\Delta} \frac{d^2\sigma \left(e^+e^- \rightarrow (c\bar{c})_{\text{singlet}}^{S_1} + (c\bar{c})_{\text{singlet}}^{S_2} \right)}{dm_{c\bar{c}_1} dm_{c\bar{c}_2}} dm_{c\bar{c}_1} dm_{c\bar{c}_2} \Bigg|_{\substack{S_1=0, S_2=0 \\ \Delta=0.5 \text{ GeV}}} = 1.2 \text{ fb}, \quad (13)$$

$$\iint_{2m_c}^{2m_D+\Delta} \frac{d^2\sigma \left(e^+e^- \rightarrow (c\bar{c})_{\text{singlet}}^{S_1} + (c\bar{c})_{\text{singlet}}^{S_2} \right)}{dm_{c\bar{c}_1} dm_{c\bar{c}_2}} dm_{c\bar{c}_1} dm_{c\bar{c}_2} \Bigg|_{\substack{S_1=0, S_2=1 \\ \Delta=0.5 \text{ GeV}}} = 19.4 \text{ fb}, \quad (14)$$

$$\iint_{2m_c}^{2m_D+\Delta} \frac{d^2\sigma \left(e^+e^- \rightarrow (c\bar{c})_{\text{singlet}}^{S_1} + (c\bar{c})_{\text{singlet}}^{S_2} \right)}{dm_{c\bar{c}_1} dm_{c\bar{c}_2}} dm_{c\bar{c}_1} dm_{c\bar{c}_2} \Bigg|_{\substack{S_1=1, S_2=1 \\ \Delta=0.5 \text{ GeV}}} = 19.8 \text{ fb}. \quad (15)$$

One can suppose that if $S_1 = 0$ and $S_2 = 1$ (or, $S_1 = 1$ and $S_2 = 0$), then J/ψ and η_c are produced in most cases; if $S_1 = 1$ and $S_2 = 1$, then J/ψ and P -wave state of quarkonium are most probably produced. Then the cross section of process

$$e^+e^- \rightarrow J/\psi + \text{charmonium} \quad (16)$$

at $\sqrt{s} = 10.6 \text{ GeV}$ can be estimated to be about 40 fb. This value can be compared with the experimental data on the charmonium pair production:

$$\sigma(e^+e^- \rightarrow J/\psi + \text{charmonium}) = 55 \pm 10 \text{ fb} \quad (\text{BELLE Collaboration}), \quad (17)$$

$$\sigma(e^+e^- \rightarrow J/\psi + \text{charmonium}) = 44.3 \pm 9 \text{ fb} \quad (\text{BaBar Collaboration}). \quad (18)$$

One can conclude that the quark-hadron duality prediction does not contradict the experimental data within experimental errors and uncertainties in m_c and α_s values. There is certainly no reason to talk about the discrepancy of order of magnitude, as it is a case for the predictions of the δ -approximation [1]. It is worth to note, that the calculation results (14) and (15) indicate a large yield of P -wave states, which is comparable with the yield of η_c and η'_c . It is also in agreement with the experimental data of the BELLE and BaBar Collaborations.

In the work [12] the quark-hadron duality hypothesis was combined with the Colour Evaporation Model predicting a yield of charmonium state with the total angular momentum J is proportional to $2J + 1$. Obviously, this model cannot be applied to the process

$e^+e^- \rightarrow J/\psi + X$ because the quantum numbers of J/ψ and X channel are strongly correlated.

The above estimations allow to predict the cross section of process $e^+e^- \rightarrow J/\psi + D\bar{D}$. Subtracting the cross section value of exclusive pair production process from the cross section value of inclusive charmonium production process (3) one can obtain:

$$\sigma(e^+e^- \rightarrow J/\psi + D + \bar{D}) \simeq 240 \text{ fb.} \quad (19)$$

That value can be compared with that obtained by the BELLE Collaboration [13]:

$$\sigma(e^+e^- \rightarrow J/\psi + c + \bar{c}) = 870 \pm 26 \text{ fb.} \quad (20)$$

Thus, our predictions underestimates the experimental results by several times. Moreover, within the model under discussion the cross section value for the process $e^+e^- \rightarrow J/\psi + D\bar{D}$ cannot be larger than the cross section value of four charmed quark production, i.e. it can be not large then 330 fb, and the essential fraction of this value should include doubly charmed baryon contribution. It is discussed in the next chapter.

III. DOUBLY CHARMED BARYON PRODUCTION

Let us estimate how often a cc -pair is produced in the color antitriplet state with a small invariant mass in a system of four c -quarks. Such an object can be considered as a candidate to create a doubly charmed baryon (see also [10]). In the duality interval $2m_c < m_{cc} < 2m_D + \Delta$ for $\alpha_s = 0.24$, $m_c = 1.25$ GeV, and $\Delta = 0.5$ GeV, one obtains the following cross section value:

$$\sigma(e^+e^- \rightarrow (cc)_{\bar{3}}/(\bar{c}\bar{c})_3 + X) = 170 \text{ fb.} \quad (21)$$

For the production of both cc - and $\bar{c}\bar{c}$ -pairs in the above duality interval, the cross section value is

$$\sigma(e^+e^- \rightarrow (cc)_{\bar{3}} + (\bar{c}\bar{c})_3) = 32 \text{ fb.} \quad (22)$$

We assume that in the kinematical region under discussion a doubly charmed baryon and an antibaryon are created with high probability. The rest of the cross section value $\sigma(e^+e^- \rightarrow (cc)_{\bar{3}}/(\bar{c}\bar{c})_3 + X)$ (about 140 fb) should be considered as an upper limit for the cross section of the processes like $e^+e^- \rightarrow \Xi_{cc} + \bar{\Lambda}_c + \bar{D}$.

Let us discuss here a relation between the inclusive charmonium production and the production of doubly charmed baryons. The c -quark and $\bar{c}$ -quark produced in the intermediate gluon splitting have rather often almost the same velocities and this provides the minimum virtuality of intermediate gluon ($\sim 4m_c^2$). Let us suppose that the color singlet $c\bar{c}$ -pair is produced with a small invariant mass. Either c -quark or $\bar{c}$ -quark in this pair is produced by the splitting gluon. For the sake of simplicity, let us suppose that it is a $\bar{c}$ -quark. Then the remaining c -quark (which does belong to the pair under consideration) is also produced by the splitting gluon. It means that there is a large probability for this c -quark to have a velocity close to the velocity of the color singlet $c\bar{c}$ -pair. This is why the remaining c -quark can create with c -quark from $c\bar{c}$ -pair an antitriplet state with a small invariant mass. Therefore such a $cc\bar{c}$ -system can create a charmonium, as well as a doubly charmed baryon. Within the δ -approximation, the relative inclusive yield of charmonia and doubly charmed baryons are determined by the ratio of squared wave functions at origin for charmonium and doubly charmed diquark, correspondingly. Unfortunately, the quark-hadron duality technique does not allow to definitely separate these yields for the case of inclusive production.

Contrary, the exclusive process of charmonium pair production and the process of pair production of doubly charmed baryons are well separated kinematically, and the problem under discussion does not arise in this case.

IV. PROCESS $e^+e^- \rightarrow J/\psi gg$

When the charmonia production in e^+e^- -annihilation is discussed, the process

$$e^+e^- \rightarrow J/\psi gg \tag{23}$$

cannot be ignored (see Fig. 2). The detailed study of the process is worthy of a separate article, because the situation here is quite similar to one with the exclusive pair production of charmonia (16). The treating of the process (16) within the δ -approximation leads to large fixed virtualities of intermediate quarks, as in the case of process (16). This means, that the δ -approximation leads to the underestimated value of the cross section of the process (23), as well as (16). The value of effective virtuality predicted within the quark-hadron duality is less than the value predicted within the δ -approximation, and it leads to the cross section value increase. In this work we do not intend to study the process $e^+e^- \rightarrow J/\psi gg$

in details and only give the cross section value predicted in the framework of hypothesis of the quark-hadron duality. For $\Delta = 0.5$ GeV and for the invariant mass of two final gluons larger than two masses of π -meson ($m_{gg} > 2M_\pi$), one can obtain:

$$\int_{2m_c}^{2m_D+\Delta} \frac{d\sigma(e^+e^- \rightarrow (c\bar{c})_{\text{singlet}}^S + gg)}{dm_{c\bar{c}}} dm_{c\bar{c}} \Big|_{\Delta=0.5 \text{ GeV}, m_{gg}>2M_\pi}^{S=1} = 1.96 \text{ pb}, \quad (24)$$

$$\int_{2m_c}^{2m_D+\Delta} \frac{d\sigma(e^+e^- \rightarrow (c\bar{c})_{\text{singlet}}^S + gg)}{dm_{c\bar{c}}} dm_{c\bar{c}} \Big|_{\Delta=0.5 \text{ GeV}, m_{gg}>2M_\pi}^{S=0} = 1.00 \text{ pb}. \quad (25)$$

If one suppose that the pair with a spin value of $S = 1$ most probably creates J/ψ -state, then the results (24) should be considered as the cross section value estimation for the process $e^+e^- \rightarrow J/\psi gg$. Of cause, this estimation is quite rough. In addition, our calculations show that the prediction value depends quite strongly on Δ . For example, at $\Delta = 0.1$ GeV the cross section value of the $e^+e^- \rightarrow J/\psi gg$ is about 0.6 pb, which is in agreement with the experimental data. To achieve a better reliability of the predictions, one can introduce an effective mass of gluon into calculations, as it was done in the works [14, 15]. Therefore, the quark-hadron duality hypothesis does not contradict the experimental data. Contrary, the predictions of the δ -approximation [15, 16]³ are too small to describe the data.

V. CONCLUSIONS

Until recently, it was believed that the experimental data on J/ψ production in e^+e^- -annihilation cannot be described within the perturbative QCD. This conclusion was based on calculation which does not take into account the motion of the valence quark inside the J/ψ -meson. Taking into account the "internal motion" by means of light cone wave function, as it is done in works [4], or by means of quark-hadron duality, as it is done in the present work, allows to resolve some contradictions between the pQCD and the experimental data. One can certainly claim that the experimental data on the charmonium pair production in e^+e^- -annihilation, as well as the experimental data on charmonium production in the process $e^+e^- \rightarrow J/\psi gg$, do not contradict pQCD predictions.

Nevertheless, the experimental value of the inclusive cross section of J/ψ production in the process $e^+e^- \rightarrow J/\psi + D + \bar{D}$ is still underestimates by theoretical predictions. Even our

³ In the work [15] the factor of 1/2 has been missed in the cross section calculation for the process $e^+e^- \rightarrow J/\psi gg$. The revised prediction value is about 0.35 pb instead of 0.7 pb.

very optimistic estimation of the cross section value (330 fb at 10.6 GeV) is approximately three times less than the experimental result. Moreover, we think that the value of about 200 fb seems to be the more reasonable estimation of the process cross section.

In conclusion it is worth to mention one more time, that the calculations done in this work within the quark-hadron duality hypothesis suggest, that the cross section of the pair production of doubly charmed baryons in e^+e^- -annihilation could be of the same order as the cross section of charmonium pair production.

We thank L. K. Gladilin and S. P. Baranov for the fruitful discussion. This work was partially supported by Russian Fund of Basic Research (project No 04-02-17530), and the CRDF (project No. M0-011-0) and was performed within the program for Support of Leading Scientific Schools (project No. 1303.2003.2). The work of A.V. Berezhnoy was also supported in part by Russian Fund of Basic Research (project No. 05-07-90292), by President of Russian Federation (project for Young PhD Support No. 2773.2005.2), and by Dynasty Foundation (project for Young Scientist Support).

-
- [1] K.-Y. Liu, Z.-G. He, and K.-T. Chao, Phys. Lett. B **557**, 45 (2003); E. Braaten and J. Lee, Phys. Rev. D **67**, 054007 (2003), Erratum-ibid. D **72**, 099901 (2005); S. P. Baranov, Phys. Atom. Nucl. **67**, 1010 (2004); V. V. Kiselev, Int. J. Mod. Phys. A **10**, 465 (1995).
 - [2] K. Abe *et al.* [Belle Collaboration], Phys. Rev. D **70**, 071102 (2004).
 - [3] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **72**, 031101(2005).
 - [4] A. E. Bondar and V. L. Chernyak, Phys. Lett. B **612**, 215 (2005); J. P. Ma and Z. G. Si, Phys. Rev. D **70**, 074007 (2004); V. V. Braguta, A. K. Likhoded, A. V. Luchinsky, Phys. Rev. D **72**, 074019 (2005), hep-ph/0602047, hep-ph/0602232.
 - [5] V. G. Kartvelishvili, A. K. Likhoded and V.A. Petrov, Phys. Lett. B **78**, 615 (1978); A. K. Likhoded, S. R. Slabospitsky, A. N. Tolstenkov, Sov. J. Nucl. Phys. **35**, 726 (1982), Yad. Fiz. **35**, 1240 (1982).
 - [6] P. Cho and A. K. Leibovich, Phys. Rev. **D54**, 6690 (1996).
 - [7] S. Baek, P. Ko, J. Lee, and H. S. Song, J.Korean Phys. Soc. **33**, 97 (1998); Phys. Rev. D **55**, 6839 (1997); Phys. Lett. B **389**, 609 (1996); K.-Y. Liu, Z.G. He, and K.-T. Chao, Phys. Rev. D **69**, 094027 (2004).

- [8] M. A. Gomshi Nobary and B. Javadi, Eur. Phys. J. C **42**, 37 (2005).
- [9] V. V. Kiselev, A. K. Likhoded, M. V. Shevlyagin, Phys. Atom. Nucl. **57**, 689 (1994);
A. K. Likhoded, A. I. Onishchenko, Phys. Atom. Nucl. **60**, 623 (1997).
- [10] V. V. Kiselev, A. K. Likhoded, and M. V. Shevlyagin, Phys. Lett. B **332**, 411 (1994);
V. V. Kiselev, A. K. Likhoded, Phys. Usp. **45**, 455 (2002).
- [11] K.Y. Liu, Z.-G. He, and K.-T. Chao, Phys. Rev. D **68** 031501 (2003); A. V. Luchinsky, Phys.
Atom. Nucl. **68**, 1400 (2005).
- [12] D. Kang, J.-W. Lee, J. Lee, T. Kim, and P. Ko, Phys. Rev. D **71**, 094019 (2005).
- [13] K. Abe *et al.* [Belle Collaboration], Phys. Rev. Lett. **88**, 052001 (2002), Phys. Rev. Lett.
89, 142001 (2002).
- [14] J. H. Field, Phys. Rev. D **66**, 013013 (2002).
- [15] A. V. Berezhnoy, A.K. Likhoded, Phys. Atom. Nucl. **67**, 757 (2004).
- [16] J. H. Kuhn and H. Schneider, Z. Phys. C **11**, 263 (1981), Phys. Rev. D **24**, 2996 (1981);
V. M. Driesen and J. H. Kuhn, Phys. Rev. D **49**, 3197 (1994); W.-Y. Keung, Phys. Rev. D
23, 2072 (1981); F. Yuan, C.-F. Qiao, and K.-T. Chao, Phys. Rev. D **56**, 321 (1997), **56**, 1663
(1997); E. Braaten and Y.-Q. Chen, Phys. Rev. Lett. **76**, 730 (1996).




Figure 1: The leading order diagrams for the process $e^+e^- \rightarrow J/\psi + c\bar{c}$. Q is momentum of the photon and q is momentum of the intermediate quark or gluon. The average of q^2 in the total phase space is about several m_c^2 . If both $c\bar{c}$ -pairs are fused into charmonia ($e^-e^+ \rightarrow J/\psi + \text{charmonium}$), then $q^2 \sim Q^2/4$ within the δ -approximation.




Figure 2: The leading order diagrams for the process $e^- e^+ \rightarrow J/\psi gg$.